

MASSIVE: Massive stAr aSteroSeIsmology, Variability, and Evolution with K2: Binary stars in Fields 2 & 3

The proposing consortium consists of scientists from Leuven University (B), Paris-Meudon Observatory (F), Liège University (B), CEA-Saclay Site (F), and Royal Observatory of Belgium, all within KASC WG3: Conny Aerts, Evelyne Alecian, Maryline Briquet, Jonas Debosscher, Peter De Cat, Pieter Degroote, Pablo Marcos-Arenal, Stéphane Mathis, Ehsan Moravveji, Coralie Neiner, Péter Pápics, Anne Thoul, Andrew Tkachenko, Santiago Andres Triana, Konstanze Zwintz

Our goal is to perform in-depth ensemble asteroseismology and variability studies of the most massive stars, with the aim to cover the full evolutionary path from the birthline to the supernova explosion. While the nominal *Kepler* mission already implied a revolution in stellar physics for solar-type stars and red giants, it was not possible to perform high-precision studies of massive OB stars or of pre-main sequence (pre-MS) stars because such targets were not sufficiently available in *Kepler's* original FoV, while CoRoT only observed a few of them, several of which during less than one month. We shall remedy this lack of data for the metal factories of the Universe, for which stellar evolution theory is least adequate while its impact on life cycles and on chemical enrichment of galaxies is dominant. The science cases that we shall address were already extensively described in the white paper by Aerts et al. (2013, arXiv:1309.3042) taking the young open cluster NGC 2244 as a case study, but this cluster cannot be observed due to the restriction of the pointing of K2 to the ecliptic. Instead, we seek to observe stars in the fields of K2 to meet the same aims but for various metallicities. This requires that we consider different classes of stars to cover the entire evolutionary path. For each sub-class of stars, we recall briefly the science case in 7 sub-proposals, including the target list for each of them.

Based on the experience of Aerts' and Neiner's teams, who were responsible for the CoRoT OB star target selection, ground-based follow-up and CoRoT data exploitation (cf. ADS since 2009), we have carefully selected the best K2 targets for our aims, as summarized in the Table below for Fields 2 & 3. Each of the targets was assigned a priority according to its rarity and expected S/N following simulations with our software (Marcos-Arenal et al., 2014, A&A in press (arXiv:1404.1886)). For these particular Fields, all our targets have highest priority. We plan to adopt the same strategy for all future K2 fields until we have light curves of sufficient quality for at least 100 members in each sub-class, to guarantee that we can place the stars in evolutionary sequences, for various masses and metallicities. For the rare objects, we request all accessible stars. Spectroscopic and spectro-polarimetric follow-up will be performed with the NARVAL, ESPADONS, and HERMES instruments for the stars brighter than 11; for fainter targets we shall apply for competitive time at ESO/IAC/OHP, where the MASSIVE consortium has high success rates. The lead PIs indicated per sub-class are members of KASC WG3, while Alecian, Debosscher, De Cat, Degroote, Marcos-Arenal, Mathis, Thoul, and Triana deliver expertise in magnetism as well as in data and theoretical modelling. The MASSIVE consortium has large expertise in analysing *Kepler* and CoRoT data.

Sub-class	PI	Field 2	Field 3	Sub-class	PI	Field 2	Field 3
Be stars	Neiner	1	0	O stars	Aerts	0	0
magnetic stars	Briquet	5	0	single B stars	Pápics	34	6
pre-MS stars	Zwintz	15	0	binary OB stars	Tkachenko	10	1
OB supergiants	Moravveji	14	3				

Binary stars are a prime source of model-independent stellar fundamental parameters, in particular of stellar masses and radii. As such, binary stars are irreplaceable for probing models of stellar structure and evolution. Recent investigations of the massive binary V380 Cyg based on *Kepler* space-photometry and ground-based spectroscopy showed that present-day single-star evolutionary models are inadequate and lack a serious amount of near-core mixing when applied to binaries (Tkachenko et al. 2014, MNRAS, 438, 3093). A larger sample of (preferably pulsating) binaries with precise masses is required to draw any firm conclusions, however. So far, for about a hundred of massive binaries only, the masses were derived to the required precision (better than 3%), as the analysis greatly relied on ground-based data (Torres et al. 2010, A&ARv, 18, 67). Moreover, the current sample is greatly biased towards short orbital period binaries (<10 d). With K2 we will significantly extend the sample of massive binary stars, in particular towards longer orbital periods. We expect the high-precision of the obtained data to provide the detection of self- and tidally-excited stellar oscillations in a significant fraction of the proposed targets. Critical information on the near-core mixing and internal rotation profile will be provided through the detailed asteroseismic analysis of these stars. Observing in long-cadence mode is sufficient for our targets.